

CERTIFIED COPY OF
PRIORITY DOCUMENT

BEST AVAILABLE COPY

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In the matter of an application for a German Patent in the name of Merck Patent GmbH, filed under No. 102 40 026.1 on 27 August 2002, and in the matter of an application for an Australian Patent.

I, Dr. Ashwood Stephen DRANE, B.Sc., Ph.D., BDÜ, translator to Steve Drane Translations Ltd. of Beechwood, Chivery, Tring, Hertfordshire, England, do solemnly and sincerely declare:


1. That I am a citizen of the United Kingdom of Great Britain and Northern Ireland.
2. That I am well acquainted with the German and English languages and am a competent translator thereof.
3. That the following is to the best of my knowledge and belief a true and correct translation of the above-referenced patent application and the Official Certificate attached thereto

Dated this 14th day of March 2007



A handwritten signature in cursive script, appearing to read 'Drane', written over a horizontal line.

Dr. Ashwood Stephen Drane



FEDERAL REPUBLIC OF GERMANY



Priority certificate regarding the filing of a patent application

File reference: 102 40 026.1

Date of filing: 27 August 2002

Applicant/proprietor: Merck Patent GmbH,
Darmstadt/DE

Title: Process for the preparation of monoalkylaminoketones

IPC: C 07 B, C 07 C, C 07 D

The attached pages are a correct and accurate reproduction of the original documents of this patent application.

Seal

Munich, 08 May 2003
German Patent and Trademark Office
On behalf of
The President

[signature]


Wehner

**Merck Patent Gesellschaft
mit beschränkter Haftung
64271 Darmstadt**

**Process for the preparation of
monoalkylaminoketones**

Process for the preparation of monoalkylaminoketones

The invention relates to processes for the preparation of monoalkylamino-ketones of the formula I




10 in which

R^1 denotes a saturated, unsaturated or aromatic carbocyclic or heterocyclic radical which is unsubstituted or mono- or polysubstituted by R^3 and/or R^4 ,

R^2 denotes alkyl having 1-20 C atoms,

15 R^3, R^4 each, independently of one another, denote H, alkyl or alkoxy having 1-20 C atoms, aryl, aryloxy or COOR^2 , F, Cl, Br, OH, CN, NO_2 , $\text{N(R}^2)_2$ or NHCOR_2 ,

by reaction of compounds of the formula II



25 in which

R^1 and R^2 have the meaning indicated above, in the presence of an alkyl-amine of the formula R^2NH_2 , in which R^2 has the meaning indicated above.

30 The compounds of the formula II are preferably employed as acid-addition salts, where, in particular, the acid-addition salts of strong acids, such as, for example, hydrohalic acid, methyl-, p-toluene- or benzenesulfonic acid, perchloric, sulfuric or phosphoric acid, are suitable. Particular preference is given to the hydrochlorides of the compounds of the formula II. On use of the acid-addition salts of the compounds of the formula II, the acid-addi-


35 tion salts of the compounds of the formula I are obtained, from which the

free bases can be liberated by addition of a strong base, such as alkali metal carbonate or hydroxide.

5 The invention facilitates, in particular, the synthesis of precursors of optically active β -monoalkylaminopropanols which are suitable as starting compounds in the preparation of medicaments, such as, for example, anti-depressants.


10 In particular, it opens up the possibility of obtaining in a simple manner 3-methylamino-1-(2-thienyl)-1-propanone, which can be used for the preparation of (S)-3-methylamino-1-(2-thienyl)-1-propanol. It is likewise possible to obtain 3-methylamino-1-phenyl-1-propanone, from which (S)-3-methylamino-1-phenyl-1-propanol can be obtained. These propanols can be, in particular, converted further, for example, into fluoxetine, tomoxetine and LY227942 (W. J. Wheeler, F. Kuo, J. Labelled Compd. Radiopharm. 15 1995, 36, 213-223).

In general, the synthesis of secondary amino ketones of the formula I under the conditions of a Mannich reaction (C. Mannich, G. Heilner, Chem. 20 Ber. 1922, 55, 362-365) from compounds of the formula III and an alkylamine of the formula R^2NH_2 in the presence of a formaldehyde source, such as paraformaldehyde, acetals of formaldehyde, such as, for example, methyl or ethyl acetals, or trioxane, proves to be difficult since the secondary amino ketone of the formula I formed primarily serves directly as 25 starting material for a subsequent second aminomethylation, where the main product obtained is the compound of the formula II:



35 In particular, this applies to the reaction of acetylthiophene IIIa with methylammonium chloride in the presence of paraformaldehyde, which

gives exclusively the dimer IIa and not the desired monomer Ia (F. F. Blicke, J. H. Burckhalter, J. Am. Chem. Soc. 1942, 64, 451-454):



The invention was therefore based on the object of finding a process for the preparation of the compounds of the formula I or salts thereof and in particular of the compound Ia or salts thereof, which can be used, in particular, as intermediates in the synthesis of medicaments, which does not have the above-mentioned disadvantages.

It has been found that the compounds of the formula I and salts thereof, which are important intermediates for the preparation of medicaments, in particular of those which exhibit, for example, actions on the central nervous system, can be obtained by reaction of compounds of the formula II or salts thereof, in particular of compounds of the formula IIa or salts thereof, in the presence of an alkylamine of the formula R²NH₂.

Above and below, the radicals R¹, R², R³, R⁴ have the meanings indicated for the formulae I to II, unless expressly stated otherwise.

In the above formulae, alkyl has 1 to 20, preferably 1 to 6, in particular 1, 2, 3 or 4 C atoms. Alkyl preferably denotes methyl or ethyl, furthermore propyl, isopropyl, furthermore also butyl, isobutyl, sec-butyl or tert-butyl.

R¹ is preferably an aromatic carbocyclic or heterocyclic radical which is unsubstituted or substituted by R³ and/or R⁴. This radical may be mono- or polycyclic and is preferably mono- or bicyclic, but in particular monocyclic. R¹ is particularly preferably unsubstituted.

If R¹ denotes a carbocyclic radical, this radical is preferably, for example, phenyl, o-, m- or p-tolyl, o-, m- or p-hydroxyphenyl, o-, m- or p-methoxyphenyl, o-, m- or p-fluorophenyl.

- 5 If R¹ denotes a heterocyclic radical, 2- or 3-furyl, 2- or 3-thienyl, 1-, 2- or 3-pyrrolyl, 1-, 2, 4- or 5-imidazolyl, 1-, 3-, 4- or 5-pyrazolyl, 2-, 4- or 5-oxazolyl, 3-, 4- or 5-isoxazolyl, 2-, 4- or 5-thiazolyl, 3-, 4- or 5-isothiazolyl, 2-, 3- or 4-pyridyl, 2-, 4-, 5- or 6-pyrimidinyl, furthermore preferably 1,2,3-triazol-1-, -4- or -5-yl, 1,2,4-triazol-1-, -3- or 5-yl, 1- or 5-tetrazolyl, 1,2,3-oxadiazol-4- or -5-yl, 1,2,4-oxadiazol-3- or -5-yl, 1,3,4-thiadiazol-2- or -5-yl, 1,2,4-thiadiazol-3- or -5-yl, 1,2,3-thiadiazol-4- or -5-yl, 3- or 4-pyridazinyl, pyrazinyl, 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl, 4- or 5-isoindolyl, 1-, 2-, 4- or 5-benzimidazolyl, 1-, 3-, 4-, 5-, 6- or 7-benzopyrazolyl, 2-, 4-, 5-, 6- or 7-benzoxazolyl, 3-, 4-, 5-, 6- or 7-benzisoxazolyl, 2-, 4-, 5-, 6- or 7-benzothiazolyl, 2-, 4-, 5-, 6- or 7-benzisothiazolyl, 4-, 5-, 6- or 7-benz-2,1,3-oxadiazolyl, 2-, 3-, 4-, 5-, 6-, 7- or 8-quinolyl, 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl, 3-, 4-, 5-, 6-, 7- or 8-cinnolyl, 2-, 4-, 5-, 6-, 7- or 8-quinazolinyl, 5- or 6-quinoxalyl, 2-, 3-, 5-, 6-, 7- or 8-2H-benzo[1,4]oxazinyl, further preferably 1,3-benzodioxol-5-yl, 1,4-benzodioxan-6-yl, 2,1,3-benzothiadiazol-4- or -5-yl or 2,1,3-benzoxadiazol-5-yl, for example, is preferably suitable. It is likewise possible to use metallocenes, such as, for example, ferrocenes, in particular acetylferrocene.

- 25 The heterocyclic radicals may also be partially or fully hydrogenated. The heterocyclic radical used can thus also be, for example, 2,3-dihydro-2-, -3-, -4- or -5-furyl, 2,5-dihydro-2-, -3-, -4- or 5-furyl, tetrahydro-2- or -3-furyl, 1,3-dioxolan-4-yl, tetrahydro-2- or -3-thienyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 2,5-dihydro-1-, -2-, -3-, -4- or -5-pyrrolyl, 1-, 2- or 3-pyrrolidinyl, tetrahydro-1-, -2- or -4-imidazolyl, 2,3-dihydro-1-, -2-, -3-, -4- or -5-pyrazolyl, tetrahydro-1-, -3- or -4-pyrazolyl, 1,4-dihydro-1-, -2-, -3- or -4-pyridyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5- or -6-pyridyl, 1-, 2-, 3- or 4-piperidinyl, 2-, 3- or 4-morpholinyl, tetrahydro-2-, -3- or -4-pyranyl, 1,4-dioxanyl, 1,3-dioxan-2-, -4- or -5-yl, hexahydro-1-, -3- or -4-pyridazinyl, hexahydro-1-, -2-, -4- or -5-pyrimidinyl, 1-, 2- or 3-piperazinyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-quinolyl, 1,2,3,4-tetrahydro-1-, -2-, -3-, -4-, -5-, -6-, -7- or -8-isoquinolyl, 2-, 3-, 5-, 6-, 7- or 8-3,4-

5 dihydro-2H-benzo[1,4]oxaziny, furthermore preferably 2,3-methylene-dioxyphenyl, 3,4-methylenedioxyphenyl, 2,3-ethylenedioxyphenyl, 3,4-ethylenedioxyphenyl, 3,4-(difluoromethylenedioxy)phenyl, 2,3-dihydro-benzofuran-5- or 6-yl, 2,3-(2-oxomethylenedioxy)phenyl or also 3,4-dihydro-2H-1,5-benzodioxepin-6- or -7-yl, furthermore preferably 2,3-dihydrobenzofuranyl or 2,3-dihydro-2-oxofuranyl.

10 The said heterocyclic radicals may additionally be substituted by R^3 and/or R^4 .

R^1 particularly preferably denotes phenyl or 2-thienyl.

15 R^2 preferably denotes methyl, ethyl, n-propyl or isopropyl, but in particular methyl.

20 R^3 and R^4 , independently of one another, preferably denote H, methyl, in particular H.

Aryloxy preferably denotes, for example, phenoxy, o-, m- or p-tolyloxy, o-, m- or p-hydroxyphenoxy, o-, m- or p-methoxyphenoxy, o-, m- or p-fluorophenoxy.

25 Aryl preferably denotes, for example, phenyl, o-, m- or p-tolyl, o-, m- or p-hydroxyphenyl, o-, m- or p-methoxyphenyl, o-, m- or p-fluorophenyl.

30 The process according to the invention is simple, with the compound of the formula II preferably being dissolved or suspended in a solvent, such as, for example, water, alcohol, ether, saturated or aromatic halogenated or halogen-free hydrocarbons or mixtures thereof. The mixture is strongly acidified by addition of a strong acid, such as, for example, hydrochloric acid or sulfuric acid. A corresponding acid-addition salt of the alkylamine of the formula R^2NH_2 can optionally also be added to the solution or suspension of the compounds of the formula II.

35 The pH of the solution is subsequently increased to about pH 2-7.5, preferably pH 4 – 7, in particular pH 5.2 to 6.8, by addition of an alkylamine of the formula R^2NH_2 , and the reaction mixture is warmed for a further 1 to


24 h, preferably 5 – 10 h, at 0° to 200°C, preferably at 10°C-100°C and in particular at 30°C – 90°C, giving the compounds of the formula I or salts thereof.

- 5 Particular preference is given to a one-pot process for the preparation of the compounds of the formula I, in which firstly the compound of the formula II is prepared by known processes, in particular in accordance with F. F. Blicke, J. H. Burckhalter, J. Am. Chem. Soc. 1942, 64, 451-454. In this process, a mixture of a formaldehyde source, such as, for example, paraformaldehyde or trioxane, is preferably reacted with a corresponding alkylammonium salt of the formula $R^2NH_2 \cdot HX$, in which HX stands for a strong acid, such as, for example, hydrogen halide, in particular hydrogen chloride, or sulfuric acid, with a ketone of the formula III and an excess of strong acid, such as, for example, hydrogen chloride, preferably in a solvent, such as, for example, water, alcohol or mixtures thereof. The reaction time of this reaction, depending on the conditions used, is generally between a few hours and 14 days, the reaction temperature is between 0°C and 200°C, normally between 10°C and 130°C, preferably between 20°C and 100°C and in particular between 30°C and 90°C. The compounds of the formula II generally precipitate from the reaction mixture as a solid after the reaction.
- 10
- 15
- 20
- 25 The pH of the hitherto strongly acidic reaction mixture comprising the compounds of the formula II is subsequently increased to about pH 2-7.5, preferably pH 5 - 6, without further isolation of this compound by addition of an alkylamine of the formula R^2NH_2 , and the reaction mixture is warmed for a further 1 to 24 h, preferably 5 – 10 h, at 0° to 200°C, preferably at 10°C-100°C and in particular at 30°C – 90°C, giving the compounds of the formula I. At high temperatures, the reaction is preferably carried out under superatmospheric pressure, preferably between 1 and 50 bar, in particular between 2 and 10 bar.
- 30

A suitable formaldehyde source is, in particular, trioxane.

- 35 A possible reaction mechanism is described below: firstly, the compound II is converted by thermal treatment into the vinyl ketone of the formula IV

- 7 -



5 and the desired hydrochloride of the compound of the formula I. Owing to the presence of methylamine, the conversion of the vinyl ketone of the formula IV into the compound of the formula I takes place simultaneously "in situ", and the latter reacts again to give the desired hydrochloride of the compound of the formula I and the vinyl ketone of the formula IV.

10 In this manner, the compound of the formula II reacts approximately completely to give the desired product of the formula I, which can be isolated comfortably after re-acidification of the reaction mixture using, for example, conc. hydrochloric acid.

15 Suitable acids for the process according to the invention are, in particular, inorganic acids, preferably non-oxidising inorganic acids.

Preferred embodiments of the process according to the invention are mentioned below:

20 Process for the preparation of compounds of the formula I, characterised in that the pH for the conversion of the compounds of the formula II into the compounds of the formula I is adjusted to about pH 2-7.5 by addition of an alkylamine of the formula R²NH₂.

25 Process for the preparation of compounds of the formula I, characterised in that the conversion of the compounds of the formula II into the compounds of the formula I is carried out at 0° - 130°C.

30 Process for the preparation of compounds of the formula I, characterised in that the conversion of the compounds of the formula II into the compounds of the formula I is carried out at 0° - 200°C, preferably under superatmospheric pressure in particular from 2 to 50 bar.

35 Process for the preparation of compounds of the formula, characterised in that firstly the compound of the formula II is obtained by reaction of a mixture of a formaldehyde source with corresponding alkylammonium salt and

a ketone of the formula III in the presence of a strong acid, and the compounds of the formula II obtained in this way are employed without further isolation for the preparation of the compounds of the formula I.

5 Process for the preparation of compounds of the formula I, characterised in that the pH of the strongly acidic reaction mixture comprising the compounds of the formula II is increased to about pH 2-7.5, without further isolation of the compound of the formula II, by addition of an alkylamine of the formula R^2NH_2 , and the mixture is subsequently warmed.

10

Process for the preparation of compounds of the formula, that the reaction mixture comprising the compounds of the formula II is warmed to 10°C to 100°C after addition of a corresponding alkylamine.

15 The process according to the invention is particularly suitable for the preparation of the ketones 3-methylamino-1-phenyl-1-propanone or 3-methylamino-1-(2-thienyl)-1-propanone, which can advantageously be converted further into the active ingredients duloxetine, fluoxetine, tomoxetine and LY227942.

20

The compounds of the formula II and also the starting materials for their preparation are, in addition, prepared by methods known per se, as described in the literature (for example in the standard works, such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart), to be precise under reaction conditions which are known and suitable for the said reactions. Use can also be made here of variants known per se which are not mentioned here in greater detail.

25

30 If desired, the starting materials can also be formed in situ by not isolating them from the reaction mixture, but instead immediately converting them further into the compounds of the formula I.

35

Some of the compounds of the formula II are known; the compounds that are not known can easily be prepared analogously to the known compounds.

Suitable solvents are, for example, hydrocarbons, such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons, such as trichloroethylene, 1,2-dichloroethane, tetrachloromethane, chloroform or dichloromethane; alcohols, such as methanol, ethanol, isopropanol, n-propanol, n-butanol or tert-butanol; ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF) or dioxane; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether, ethylene glycol dimethyl ether (diglyme), if desired also mixtures of the said solvents with one another or mixtures with water.

A base of the formula I can be converted into the associated acid-addition salt using an acid, for example by reaction of equivalent amounts of the base and the acid in an inert solvent, such as ethanol, followed by evaporation. Particularly suitable acids for this reaction are those which give physiologically acceptable salts. Thus, it is possible to use inorganic acids, for example sulfuric acid, nitric acid, hydrohalic acids, such as hydrochloric acid or hydrobromic acid, phosphoric acids, such as orthophosphoric acid, sulfamic acid, furthermore organic acids, in particular aliphatic, alicyclic, araliphatic, aromatic or heterocyclic mono- or polybasic carboxylic, sulfonic or sulfuric acids, for example formic acid, acetic acid, propionic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, citric acid, gluconic acid, ascorbic acid, nicotinic acid, isonicotinic acid, methane- or ethanesulfonic acid, ethanedisulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, naphthalenemono- and disulfonic acids, laurylsulfuric acid. Salts with physiologically unacceptable acids, for example picrates, can be used for the isolation and/or purification of the compounds of the formula I.

On the other hand, compounds of the formula I can be converted into the corresponding metal salts, in particular alkali metal or alkaline earth metal salts, or into the corresponding ammonium salts using bases (for example sodium hydroxide, sodium carbonate, potassium hydroxide or potassium carbonate).

The invention furthermore relates to the use of the compounds of the formula I as intermediates for the synthesis of medicaments. Corresponding medicaments are mentioned, for example, in Synlett, 689-690, 1991.

- 5 The invention furthermore relates to the use of the compounds of the formula I as intermediates for the synthesis of medicaments which exhibit actions on the central nervous system.

10 Above and below, all temperatures are indicated in °C. In the following examples, "conventional work-up" means: if necessary, water is added, the pH is adjusted, if necessary, to values between 2 and 10, depending on the constitution of the end product, the mixture is extracted with ethyl acetate or dichloromethane, the phases are separated, the organic phase is dried over sodium sulfate and evaporated, and the product is purified by
15 chromatography on silica gel and/or by crystallisation. R_f values on silica gel.

Example 1:


20 A mixture of 49 g of trioxane, 111 g of methylammonium chloride, 162.2 g of acetylthiophene and 12 g of 37% hydrochloric acid in 176 ml of ethanol and 44 ml of water is refluxed for 17 h. 17.6 g of methylamine solution (40% in water) are subsequently added, and the mixture is warmed at 65-84°C for 7 h. The reaction mixture is then allowed to cool to room temperature, 23.7 g of 37% hydrochloric acid are added, and the mixture is cooled
25 to below 0°C. The deposited crystals are filtered off with suction, washed with acetone and subsequently dried, giving the desired ketone.

Example 2:

30 A mixture of 45.2 g of trioxane, 102.3 g of methylammonium chloride, 127.3 g of acetylthiophene and 10 ml of 37% hydrochloric acid in 242 ml of ethanol and 61 ml of water is refluxed for 19 h. The mixture is subsequently diluted with 400 ml of ethanol, 19.9 g of methylamine solution (40% in water) are added, and the mixture is again refluxed for 7 h. The reaction mixture is then allowed to cool firstly to room temperature and is cooled at
35 -15°C for 48 h. The deposited crystals are filtered off with suction, washed with 90 g of ethanol and subsequently dried in vacuo at 45°C for 17 h.

Patent Claims

1. Process for the preparation of monoalkylaminoketones of the formula I




in which

R^1 denotes a saturated, unsaturated or aromatic heterocyclic radical which is unsubstituted or mono- or polysubstituted by R^3 and/or R^4 ,

R^2 denotes alkyl having 1-20 C atoms,

R^3, R^4 each, independently of one another, denote H, alkyl or alkoxy having 1-20 C atoms, aryl, aryloxy or COOR^2 , F, Cl, Br, OH, CN, NO_2 , $\text{N}(\text{R}^2)_2$ or NHCOR_2 ,

by reaction of compounds of the formula II



in which


R^1 and R^2 have the meaning indicated above, in the presence of an alkylamine of the formula R^2NH_2 , in which R^2 has the meaning indicated above.

2. Process according to Claim 1, in which R^1 denotes phenyl or 2-thienyl.
3. Process according to Claim 1 or 2, in which R^2 denotes methyl, ethyl, n-propyl or isopropyl.
4. Process for the preparation of compounds of the formula I according to one or more of Claims 1 to 3, characterised in that the pH for the conversion of the compounds of the formula II into the compounds of the

formula I is adjusted to about pH 2-7.5 by addition of an alkylamine of the formula R^2NH_2 .

5. Process for the preparation of compounds of the formula I according to one or more of Claims 1 to 4, characterised in that the conversion of the compounds of the formula II into the compounds of the formula I is carried out at $0^\circ - 200^\circ C$.

6. Process for the preparation of compounds of the formula I according to one or more of Claims 1 to 5, characterised in that firstly the compound of the formula II is obtained by reaction of a mixture of a form-aldehyde source with a corresponding alkylammonium salt and a ketone of the formula III



in which R^1 has the meaning indicated in Claim 1, in the presence of a strong acid, and the compounds of the formula II obtained in this way are employed without further isolation for the preparation of the compounds of the formula I.

7. Process for the preparation of compounds of the formula I according to Claim 6, characterised in that the pH of the strongly acidic reaction mixture comprising the compounds of the formula II is increased to about pH 2-7.5, without further isolation of this compound, by addition of an alkylamine of the formula R^2NH_2 , and the mixture is subsequently warmed.

8. Process for the preparation of compounds of the formula I according to Claim 7, characterised in that the reaction mixture comprising the compounds of the formula II is warmed to $0^\circ C$ to $200^\circ C$ after addition of a corresponding alkylamine.

9. Process according to one or more of Claims 1 to 8 for the preparation of 3-methylamino-1-phenyl-1-propanone or 3-methylamino-1-(2-thienyl)-1-propanone.

10. Process according to one or more of Claims 1 to 9, characterised in
that an acid-addition salt of the compound of the formula II is
employed, and an acid-addition salt of the compound of the formula I
is obtained.

5

10

15

20


25

30

35

Abstract

The invention relates to processes for the preparation of monoalkylamino-ketones of the formula I



10 in which R¹ and R² have the meanings indicated.

15

20

25

30

35